

Energy cost of entanglement extraction from quantum fields

QISS HKU Workshop 2020

Hong Kong

Lucas Hackl

with Robert H. Jonsson (Max Planck Institute of Quantum Optics, Garching)

[[arXiv:1904.06246](#), *Quantum* 3, 165 (2019)]

ENTANGLEMENT is a hallmark of quantum theory and a resource for quantum applications:

- Super-dense coding [Bennett, Wiesner '92]
- Quantum cryptography [Bennett '92]
- Quantum teleportation [Bennett et. al. '93]

ENTANGLEMENT is a hallmark of quantum theory and a resource for quantum applications:

- Super-dense coding [Bennett, Wiesner '92]
- Quantum cryptography [Bennett '92]
- Quantum teleportation [Bennett et. al. '93]

All these applications require ENTANGLED STATES across specified system decompositions:

- Ground states of **field theories with local Hamiltonians** are *spatially highly entangled*.
⇒ Area laws of entanglement entropies! [Eisert, Cramer, Plenio '08]

ENTANGLEMENT is a hallmark of quantum theory and a resource for quantum applications:

- Super-dense coding [Bennett, Wiesner '92]
- Quantum cryptography [Bennett '92]
- Quantum teleportation [Bennett et. al. '93]

All these applications require ENTANGLED STATES across specified system decompositions:

- Ground states of **field theories with local Hamiltonians** are *spatially highly entangled*.
⇒ Area laws of entanglement entropies! [Eisert, Cramer, Plenio '08]
- Can we utilize this entanglement to make it *usable for physical applications*?
⇒ We would like to extract an amount ΔS of entanglement entropy.

ENTANGLEMENT is a hallmark of quantum theory and a resource for quantum applications:

- Super-dense coding [Bennett, Wiesner '92]
- Quantum cryptography [Bennett '92]
- Quantum teleportation [Bennett et. al. '93]

All these applications require ENTANGLED STATES across specified system decompositions:

- Ground states of **field theories with local Hamiltonians** are *spatially highly entangled*.
⇒ Area laws of entanglement entropies! [Eisert, Cramer, Plenio '08]
- Can we utilize this entanglement to make it *usable for physical applications*?
⇒ We would like to extract an amount ΔS of entanglement entropy.
- Extraction of entanglement from the ground state $|\psi\rangle$ *raises energy* [Bény et. al. '18]:
⇒ Change of state $|\psi\rangle \rightarrow |\psi'\rangle$ implies energy difference $\Delta E = \langle\psi'|\hat{H}|\psi'\rangle - \langle\psi|\hat{H}|\psi\rangle$.

ENTANGLEMENT is a hallmark of quantum theory and a resource for quantum applications:

- Super-dense coding [Bennett, Wiesner '92]
- Quantum cryptography [Bennett '92]
- Quantum teleportation [Bennett et. al. '93]

All these applications require ENTANGLED STATES across specified system decompositions:

- Ground states of **field theories with local Hamiltonians** are *spatially highly entangled*.
⇒ Area laws of entanglement entropies! [Eisert, Cramer, Plenio '08]
- Can we utilize this entanglement to make it *usable for physical applications*?
⇒ We would like to extract an amount ΔS of entanglement entropy.
- Extraction of entanglement from the ground state $|\psi\rangle$ *raises energy* [Bény et. al. '18]:
⇒ Change of state $|\psi\rangle \rightarrow |\psi'\rangle$ implies energy difference $\Delta E = \langle\psi'| \hat{H} |\psi'\rangle - \langle\psi| \hat{H} |\psi\rangle$.

Key question of this talk:

What is the minimal price in energy ΔE to extract ΔS of entanglement?


Our setup, protocol and results

Our setup

Bosonic and fermionic field theories with quadratic Hamiltonian and Gaussian ground state $|\psi\rangle$:

$$\hat{H} = \int d^3x \frac{1}{2} h_{ab} \hat{\xi}^a(x) \hat{\xi}^b(x) \quad \text{with} \quad \hat{\xi}^a(x) \text{ representing bosonic or fermionic fields}$$

Examples: Klein-Gordon field, Dirac field in Minkowski space or curved backgrounds.



Our setup

Bosonic and fermionic field theories with quadratic Hamiltonian and Gaussian ground state $|\psi\rangle$:

$$\hat{H} = \int d^3x \frac{1}{2} h_{ab} \hat{\xi}^a(x) \hat{\xi}^b(x) \quad \text{with} \quad \hat{\xi}^a(x) \text{ representing bosonic or fermionic fields}$$


Examples: Klein-Gordon field, Dirac field in Minkowski space or curved backgrounds.

- ① A chosen mode A consists of:

$$\hat{Q}_A = \int d^3x q_a(x) \hat{\xi}^a(x)$$

$$\hat{P}_A = \int d^3x p_a(x) \hat{\xi}^a(x)$$

$$\text{s. t. } [\hat{Q}_A, \hat{P}_A]_{\pm} = i$$



Our setup

Bosonic and fermionic field theories with quadratic Hamiltonian and Gaussian ground state $|\psi\rangle$:

$$\hat{H} = \int d^3x \frac{1}{2} h_{ab} \hat{\xi}^a(x) \hat{\xi}^b(x) \quad \text{with} \quad \hat{\xi}^a(x) \text{ representing bosonic or fermionic fields}$$

Examples: Klein-Gordon field, Dirac field in Minkowski space or curved backgrounds.

- ① A chosen mode A consists of:

$$\hat{Q}_A = \int d^3x q_a(x) \hat{\xi}^a(x)$$

$$\hat{P}_A = \int d^3x p_a(x) \hat{\xi}^a(x)$$


$$\text{s. t. } [\hat{Q}_A, \hat{P}_A]_{\pm} = i$$

- ② Every mode A has a partner \bar{A}

$$\hat{Q}_{\bar{A}} = \int d^3x \bar{q}_a(x) \hat{\xi}^a(x)$$

$$\hat{P}_{\bar{A}} = \int d^3x \bar{p}_a(x) \hat{\xi}^a(x)$$

$$\text{s. t. } |\psi\rangle = |\psi_{A\bar{A}}\rangle \otimes |\psi_{\text{Rest}}\rangle$$



Our setup

Bosonic and fermionic field theories with quadratic Hamiltonian and Gaussian ground state $|\psi\rangle$:

$$\hat{H} = \int d^3x \frac{1}{2} h_{ab} \hat{\xi}^a(x) \hat{\xi}^b(x) \quad \text{with} \quad \hat{\xi}^a(x) \text{ representing bosonic or fermionic fields}$$

Examples: Klein-Gordon field, Dirac field in Minkowski space or curved backgrounds.

- ① A chosen mode A consists of:

$$\hat{Q}_A = \int d^3x q_a(x) \hat{\xi}^a(x)$$

$$\hat{P}_A = \int d^3x p_a(x) \hat{\xi}^a(x)$$


$$\text{s. t. } [\hat{Q}_A, \hat{P}_A]_{\pm} = i$$

- ② Every mode A has a partner \bar{A}

$$\hat{Q}_{\bar{A}} = \int d^3x \bar{q}_a(x) \hat{\xi}^a(x)$$

$$\hat{P}_{\bar{A}} = \int d^3x \bar{p}_a(x) \hat{\xi}^a(x)$$

$$\text{s. t. } |\psi\rangle = |\psi_{A\bar{A}}\rangle \otimes |\psi_{\text{Rest}}\rangle$$




$\Rightarrow A$ and \bar{A} are entangled, but together they are in a pure state $|\psi_{A\bar{A}}\rangle$!

Entanglement extraction:

- 1 Start with $|\psi\rangle = |\psi_{A\bar{A}}\rangle \otimes |\psi_{\text{Rest}}\rangle$


Field theory model



Our protocol

Entanglement extraction:


- 1 Start with $|\psi\rangle = |\psi_{A\bar{A}}\rangle \otimes |\psi_{\text{Rest}}\rangle$
- 2 Alice couples to mode A



Our protocol

Entanglement extraction:


- 1 Start with $|\psi\rangle = |\psi_{A\bar{A}}\rangle \otimes |\psi_{\text{Rest}}\rangle$
- 2 Alice couples to mode A
- 3 Find partner mode \bar{A}



Our protocol

Entanglement extraction:


- 1 Start with $|\psi\rangle = |\psi_{A\bar{A}}\rangle \otimes |\psi_{\text{Rest}}\rangle$
- 2 Alice couples to mode A
- 3 Find partner mode \bar{A}
- 4 Bob couples to partner \bar{A}



Our protocol

Entanglement extraction:

- 1 Start with $|\psi\rangle = |\psi_{A\bar{A}}\rangle \otimes |\psi_{\text{Rest}}\rangle$
- 2 Alice couples to mode A
- 3 Find partner mode \bar{A}
- 4 Bob couples to partner \bar{A}
- 5 Swap $|\psi_{A\bar{A}}\rangle$ with $|\psi_{\text{Alice}}\rangle \otimes |\psi_{\text{Bob}}\rangle$
 $\Rightarrow |\psi'\rangle = |\psi_{\text{Alice}}\rangle \otimes |\psi_{\text{Bob}}\rangle \otimes |\psi_{\text{Rest}}\rangle$




Our protocol

Entanglement extraction:

- 1 Start with $|\psi\rangle = |\psi_{A\bar{A}}\rangle \otimes |\psi_{\text{Rest}}\rangle$
- 2 Alice couples to mode A
- 3 Find partner mode \bar{A}
- 4 Bob couples to partner \bar{A}
- 5 Swap $|\psi_{A\bar{A}}\rangle$ with $|\psi_{\text{Alice}}\rangle \otimes |\psi_{\text{Bob}}\rangle$
 $\Rightarrow |\psi'\rangle = |\psi_{\text{Alice}}\rangle \otimes |\psi_{\text{Bob}}\rangle \otimes |\psi_{\text{Rest}}\rangle$

Resulting change:




Our protocol

Entanglement extraction:

- 1 Start with $|\psi\rangle = |\psi_{A\bar{A}}\rangle \otimes |\psi_{\text{Rest}}\rangle$
- 2 Alice couples to mode A
- 3 Find partner mode \bar{A}
- 4 Bob couples to partner \bar{A}
- 5 Swap $|\psi_{A\bar{A}}\rangle$ with $|\psi_{\text{Alice}}\rangle \otimes |\psi_{\text{Bob}}\rangle$
 $\Rightarrow |\psi'\rangle = |\psi_{\text{Alice}}\rangle \otimes |\psi_{\text{Bob}}\rangle \otimes |\psi_{\text{Rest}}\rangle$

Resulting change:

- Alice and Bob share entanglement ΔS




Our protocol

Entanglement extraction:

- 1 Start with $|\psi\rangle = |\psi_{A\bar{A}}\rangle \otimes |\psi_{\text{Rest}}\rangle$
- 2 Alice couples to mode A
- 3 Find partner mode \bar{A}
- 4 Bob couples to partner \bar{A}
- 5 Swap $|\psi_{A\bar{A}}\rangle$ with $|\psi_{\text{Alice}}\rangle \otimes |\psi_{\text{Bob}}\rangle$
 $\Rightarrow |\psi'\rangle = |\psi_{\text{Alice}}\rangle \otimes |\psi_{\text{Bob}}\rangle \otimes |\psi_{\text{Rest}}\rangle$

Resulting change:

- Alice and Bob share entanglement ΔS
- Energy is increased by ΔE



Our protocol


Entanglement extraction:

- 1 Start with $|\psi\rangle = |\psi_{A\bar{A}}\rangle \otimes |\psi_{\text{Rest}}\rangle$
- 2 Alice couples to mode A
- 3 Find partner mode \bar{A}
- 4 Bob couples to partner \bar{A}
- 5 Swap $|\psi_{A\bar{A}}\rangle$ with $|\psi_{\text{Alice}}\rangle \otimes |\psi_{\text{Bob}}\rangle$
 $\Rightarrow |\psi'\rangle = |\psi_{\text{Alice}}\rangle \otimes |\psi_{\text{Bob}}\rangle \otimes |\psi_{\text{Rest}}\rangle$

Resulting change:



- Alice and Bob share entanglement ΔS
- Energy is increased by ΔE

What is the minimal ΔE
to extract a fixed amount ΔS ?



Our result

Hamiltonian $\hat{H} = \int_k d^3k \omega_k \hat{n}_k + E_0$ has **minimal energy** cost of entanglement extraction:



- ϵ_1 and ϵ_2 are the smallest excitation energies, e.g., $\epsilon_1 = \epsilon_2 = m$ for most field theories.
- In both cases, we find $\lim_{\Delta S \rightarrow 0} \Delta E/\Delta S = 0$, i.e., extracting small amounts ΔS is cheap.
- For bosons, we have $\Delta E \sim \frac{\sqrt{\epsilon_1 \epsilon_2}}{2} \exp(\Delta S)$, while $\Delta E \leq \epsilon_1$ for fermions.

Derivation (for bosons)

① **Assume that we already solved the problem.**

We found A and \bar{A} that share entanglement ΔS with minimal energy cost ΔE .

Derivation strategy (for bosons)

- 1 **Assume that we already solved the problem.**

We found A and \bar{A} that share entanglement ΔS with minimal energy cost ΔE .

- 2 **Restrict state and Hamiltonian to subsystem $A \oplus \bar{A}$.**

Here, the state $|\psi_{A\bar{A}}\rangle$ is the ground state of the restricted Hamiltonian

$$\hat{H}_{A\bar{A}} = \underbrace{\frac{\epsilon_1}{2}(\hat{Q}_1^2 + \hat{P}_1^2)}_{\hat{H}_1} + \underbrace{\frac{\epsilon_2}{2}(\hat{Q}_2^2 + \hat{P}_2^2)}_{\hat{H}_2}$$

w.r.t. some basis $\hat{\xi}^a = (\hat{Q}_1, \hat{P}_1, \hat{Q}_2, \hat{P}_2)$. We also have $\min(\omega_k) \leq \epsilon_i \leq \max(\omega_k)$.
Note that $|\psi_{A\bar{A}}\rangle$ does not have any entanglement across 1 and 2.

Derivation strategy (for bosons)

- 1 **Assume that we already solved the problem.**

We found A and \bar{A} that share entanglement ΔS with minimal energy cost ΔE .

- 2 **Restrict state and Hamiltonian to subsystem $A \oplus \bar{A}$.**

Here, the state $|\psi_{A\bar{A}}\rangle$ is the ground state of the restricted Hamiltonian

$$\hat{H}_{A\bar{A}} = \underbrace{\frac{\epsilon_1}{2}(\hat{Q}_1^2 + \hat{P}_1^2)}_{\hat{H}_1} + \underbrace{\frac{\epsilon_2}{2}(\hat{Q}_2^2 + \hat{P}_2^2)}_{\hat{H}_2}$$

w.r.t. some basis $\hat{\xi}^a = (\hat{Q}_1, \hat{P}_1, \hat{Q}_2, \hat{P}_2)$. We also have $\min(\omega_k) \leq \epsilon_i \leq \max(\omega_k)$.

Note that $|\psi_{A\bar{A}}\rangle$ does not have any entanglement across 1 and 2.

- 3 **Assume $\epsilon_1 = \epsilon_2 = \epsilon$ to simplify argument.**

Derivation strategy (for bosons)

- 1 **Assume that we already solved the problem.**

We found A and \bar{A} that share entanglement ΔS with minimal energy cost ΔE .

- 2 **Restrict state and Hamiltonian to subsystem $A \oplus \bar{A}$.**

Here, the state $|\psi_{A\bar{A}}\rangle$ is the ground state of the restricted Hamiltonian

$$\hat{H}_{A\bar{A}} = \underbrace{\frac{\epsilon_1}{2}(\hat{Q}_1^2 + \hat{P}_1^2)}_{\hat{H}_1} + \underbrace{\frac{\epsilon_2}{2}(\hat{Q}_2^2 + \hat{P}_2^2)}_{\hat{H}_2}$$

w.r.t. some basis $\hat{\xi}^a = (\hat{Q}_1, \hat{P}_1, \hat{Q}_2, \hat{P}_2)$. We also have $\min(\omega_k) \leq \epsilon_i \leq \max(\omega_k)$.
Note that $|\psi_{A\bar{A}}\rangle$ does not have any entanglement across 1 and 2.

- 3 **Assume $\epsilon_1 = \epsilon_2 = \epsilon$ to simplify argument.**
- 4 **Optimize within the system $A \oplus \bar{A}$ and find ΔE_{\min} for fixed ΔS .**

Derivation strategy (for bosons)

- 1 **Assume that we already solved the problem.**

We found A and \bar{A} that share entanglement ΔS with minimal energy cost ΔE .

- 2 **Restrict state and Hamiltonian to subsystem $A \oplus \bar{A}$.**

Here, the state $|\psi_{A\bar{A}}\rangle$ is the ground state of the restricted Hamiltonian

$$\hat{H}_{A\bar{A}} = \underbrace{\frac{\epsilon_1}{2}(\hat{Q}_1^2 + \hat{P}_1^2)}_{\hat{H}_1} + \underbrace{\frac{\epsilon_2}{2}(\hat{Q}_2^2 + \hat{P}_2^2)}_{\hat{H}_2}$$

w.r.t. some basis $\hat{\xi}^a = (\hat{Q}_1, \hat{P}_1, \hat{Q}_2, \hat{P}_2)$. We also have $\min(\omega_k) \leq \epsilon_i \leq \max(\omega_k)$.
Note that $|\psi_{A\bar{A}}\rangle$ does not have any entanglement across 1 and 2.

- 3 **Assume $\epsilon_1 = \epsilon_2 = \epsilon$ to simplify argument.**
- 4 **Optimize within the system $A \oplus \bar{A}$ and find ΔE_{\min} for fixed ΔS .**
- 5 **Optimize within all possible choice for $A \oplus \bar{A}$.**

Explicit derivation steps (for bosons)

Let us perform the last two steps explicitly:

Explicit derivation steps (for bosons)

Let us perform the last two steps explicitly:

- ④ **Optimize within the system $A \oplus \bar{A}$ and find ΔE_{\min} for fixed ΔS**

Perform Bogoliubov transformation to find basis $(\hat{Q}_A, \hat{P}_A, \hat{Q}_{\bar{A}}, \hat{P}_{\bar{A}})$ with

$$\Delta S = s_b(\text{ch } 2r) \quad \text{with} \quad s_b(x) = \left(\frac{x+1}{2}\right) \log\left(\frac{x+1}{2}\right) - \left(\frac{x-1}{2}\right) \log\left(\frac{x-1}{2}\right)$$

Explicit derivation steps (for bosons)

Let us perform the last two steps explicitly:

- ④ **Optimize within the system $A \oplus \bar{A}$ and find ΔE_{\min} for fixed ΔS**

Perform Bogoliubov transformation to find basis $(\hat{Q}_A, \hat{P}_A, \hat{Q}_{\bar{A}}, \hat{P}_{\bar{A}})$ with

$$\Delta S = s_b(\text{ch } 2r) \quad \text{with} \quad s_b(x) = \left(\frac{x+1}{2}\right) \log\left(\frac{x+1}{2}\right) - \left(\frac{x-1}{2}\right) \log\left(\frac{x-1}{2}\right)$$

$$\hat{H}_{A\bar{A}} = \underbrace{\frac{\epsilon \text{ch } 2r}{2} (\hat{Q}_A^2 + \hat{P}_A^2)}_{\hat{H}_A} + \underbrace{\frac{\epsilon \text{ch } 2r}{2} (\hat{Q}_{\bar{A}}^2 + \hat{P}_{\bar{A}}^2)}_{\hat{H}_{\bar{A}}} - \underbrace{\epsilon \text{sh } 2r (\hat{Q}_{\bar{A}} \hat{P}_A + \hat{Q}_A \hat{P}_{\bar{A}})}_{\hat{H}_I}$$

Explicit derivation steps (for bosons)

Let us perform the last two steps explicitly:

- ④ **Optimize within the system $A \oplus \bar{A}$ and find ΔE_{\min} for fixed ΔS**

Perform Bogoliubov transformation to find basis $(\hat{Q}_A, \hat{P}_A, \hat{Q}_{\bar{A}}, \hat{P}_{\bar{A}})$ with

$$\Delta S = s_b(\text{ch } 2r) \quad \text{with} \quad s_b(x) = \left(\frac{x+1}{2}\right) \log\left(\frac{x+1}{2}\right) - \left(\frac{x-1}{2}\right) \log\left(\frac{x-1}{2}\right)$$

$$\hat{H}_{A\bar{A}} = \underbrace{\frac{\epsilon \text{ch } 2r}{2}(\hat{Q}_A^2 + \hat{P}_A^2)}_{\hat{H}_A} + \underbrace{\frac{\epsilon \text{ch } 2r}{2}(\hat{Q}_{\bar{A}}^2 + \hat{P}_{\bar{A}}^2)}_{\hat{H}_{\bar{A}}} - \underbrace{\epsilon \text{sh } 2r(\hat{Q}_{\bar{A}}\hat{P}_A + \hat{Q}_A\hat{P}_{\bar{A}})}_{\hat{H}_I}$$

$$\Delta E = \epsilon \text{ch } 2r - \epsilon = 2\epsilon \text{sh}^2 r$$

Explicit derivation steps (for bosons)

Let us perform the last two steps explicitly:

- ④ **Optimize within the system $A \oplus \bar{A}$ and find ΔE_{\min} for fixed ΔS**

Perform Bogoliubov transformation to find basis $(\hat{Q}_A, \hat{P}_A, \hat{Q}_{\bar{A}}, \hat{P}_{\bar{A}})$ with

$$\Delta S = s_b(\text{ch } 2r) \quad \text{with} \quad s_b(x) = \left(\frac{x+1}{2}\right) \log\left(\frac{x+1}{2}\right) - \left(\frac{x-1}{2}\right) \log\left(\frac{x-1}{2}\right)$$

$$\hat{H}_{A\bar{A}} = \underbrace{\frac{\epsilon \text{ch } 2r}{2} (\hat{Q}_A^2 + \hat{P}_A^2)}_{\hat{H}_A} + \underbrace{\frac{\epsilon \text{ch } 2r}{2} (\hat{Q}_{\bar{A}}^2 + \hat{P}_{\bar{A}}^2)}_{\hat{H}_{\bar{A}}} - \underbrace{\epsilon \text{sh } 2r (\hat{Q}_{\bar{A}} \hat{P}_A + \hat{Q}_A \hat{P}_{\bar{A}})}_{\hat{H}_I}$$

$$\Delta E = \epsilon \text{ch } 2r - \epsilon = 2\epsilon \text{sh}^2 r$$

$$= \frac{1}{2} \left(\sqrt{\epsilon_1^2 + \epsilon_2^2 + 2\epsilon_1\epsilon_2 \text{ch } 4r} - \epsilon_1 - \epsilon_2 \right) \quad \text{for } \epsilon_1 \neq \epsilon_2$$

Explicit derivation steps (for bosons)

Let us perform the last two steps explicitly:

- ④ **Optimize within the system $A \oplus \bar{A}$ and find ΔE_{\min} for fixed ΔS**

Perform Bogoliubov transformation to find basis $(\hat{Q}_A, \hat{P}_A, \hat{Q}_{\bar{A}}, \hat{P}_{\bar{A}})$ with

$$\Delta S = s_b(\text{ch } 2r) \quad \text{with} \quad s_b(x) = \left(\frac{x+1}{2}\right) \log\left(\frac{x+1}{2}\right) - \left(\frac{x-1}{2}\right) \log\left(\frac{x-1}{2}\right)$$

$$\hat{H}_{A\bar{A}} = \underbrace{\frac{\epsilon \text{ch } 2r}{2} (\hat{Q}_A^2 + \hat{P}_A^2)}_{\hat{H}_A} + \underbrace{\frac{\epsilon \text{ch } 2r}{2} (\hat{Q}_{\bar{A}}^2 + \hat{P}_{\bar{A}}^2)}_{\hat{H}_{\bar{A}}} - \underbrace{\epsilon \text{sh } 2r (\hat{Q}_{\bar{A}} \hat{P}_A + \hat{Q}_A \hat{P}_{\bar{A}})}_{\hat{H}_I}$$

$$\Delta E = \epsilon \text{ch } 2r - \epsilon = 2\epsilon \text{sh}^2 r$$

$$= \frac{1}{2} \left(\sqrt{\epsilon_1^2 + \epsilon_2^2 + 2\epsilon_1 \epsilon_2 \text{ch } 4r} - \epsilon_1 - \epsilon_2 \right) \quad \text{for } \epsilon_1 \neq \epsilon_2$$

- ⑤ **Optimize within all possible choice for $A \oplus \bar{A}$**

We find that ΔE is monotonic in ϵ_1 and ϵ_2 , so choosing them minimally is optimal.



For quantum fields with mass gap, this implies $\epsilon_1 = \epsilon_2 = m$.

Summary

Summary

Summary of key findings:

- Minimal energy cost of entanglement extraction in quadratic systems



- Applies to bosonic and fermionic field theories (and lattice systems)

Outlook for next steps:

- Tighten bounds by including locality assumptions for Alice and Bob.
- Develop perturbation methods to treat interacting systems.